

# Discrete Math Review

Dr. Chuck Rocca  
roccac@wcsu.edu

<http://sites.wcsu.edu/roccac>



# Table of Contents

- 1 Sets, Relations, and Functions
- 2 Graph Theory
- 3 Theorems and Proofs
- 4 Next Class



# Table of Contents

1 Sets, Relations, and Functions

2 Graph Theory

3 Theorems and Proofs

4 Next Class




# Sets

- Sets:

$$A = \{a, e, i, o, u, y\} \text{ and}$$

$$B = \{b, c, d, f, g, h, \dots, z\}$$




# Sets

- Sets:

$$A = \{a, e, i, o, u, y\} \text{ and}$$

$$B = \{b, c, d, f, g, h, \dots, z\}$$



# Sets


- Sets:

$$A = \{a, e, i, o, u, y\} \text{ and}$$

$$B = \{b, c, d, f, g, h, \dots, z\}$$

- Union:

$$A \cup B = \{a, b, c, d, e, \dots, z\}$$



# Sets

- Sets:

$$A = \{a, e, i, o, u, y\} \text{ and}$$


$$B = \{b, c, d, f, g, h, \dots, z\}$$

- Union:

$$A \cup B = \{a, b, c, d, e, \dots, z\}$$

- Intersection:

$$A \cap B = \{y\}$$



## Sets

- Sets:

$$A = \{a, e, i, o, u, y\} \text{ and}$$

$$B = \{b, c, d, f, g, h, \dots, z\}$$

- Union:


$$A \cup B = \{a, b, c, d, e, \dots, z\}$$

- Intersection:

$$A \cap B = \{y\}$$

- Complement:

$$A^c = (B \setminus \{y\}) \cup \{0, 1, \dots, 9\}$$





# Sets

- Sets:

$$A = \{a, e, i, o, u, y\} \text{ and}$$

$$B = \{b, c, d, f, g, h, \dots, z\}$$

- Union:

$$A \cup B = \{a, b, c, d, e, \dots, z\}$$

- Intersection:


$$A \cap B = \{y\}$$

- Complement:

$$A^c = (B \setminus \{y\}) \cup \{0, 1, \dots, 9\}$$

- Universal Set:

$$\mathcal{U} = A \cup B \cup \{0, 1, \dots, 9\}$$



# New Sets from Old

- $A = \{a, b, c\}$  and  $B = \{0, 1, 2\}$



# New Sets from Old

- $A = \{a, b, c\}$  and  $B = \{0, 1, 2\}$
- Cartesian Product:

$$A \times B = \{(a, 0), (a, 1), (a, 2), (b, 0), (b, 1), (b, 2), (c, 0), (c, 1), (c, 2)\}$$



# New Sets from Old

- $A = \{a, b, c\}$  and  $B = \{0, 1, 2\}$

- Cartesian Product:

$$A \times B = \{(a, 0), (a, 1), (a, 2), (b, 0), (b, 1), (b, 2), (c, 0), (c, 1), (c, 2)\}$$

- Power Set:

$$\mathcal{P}(A) = \{\emptyset, \{a\}, \{b\}, \{c\}, \{a, b\}, \{a, c\}, \{b, c\}, \{a, b, c\}\}$$

$$|\mathcal{P}(A)| = 2^{|A|}$$



# Cardinality of a Power Set: $|\mathcal{P}(S)| = 2^{|S|}$

- $A = \{0, 1\}$  and  $B = \{0, 1, 2\}$



# Cardinality of a Power Set: $|\mathcal{P}(S)| = 2^{|S|}$

- $A = \{0, 1\}$  and  $B = \{0, 1, 2\}$
- $\mathcal{P}(A) = \{\emptyset, \{0\}, \{1\}, \{0, 1\}\}$



# Cardinality of a Power Set: $|\mathcal{P}(S)| = 2^{|S|}$

- $A = \{0, 1\}$  and  $B = \{0, 1, 2\}$
- $\mathcal{P}(A) = \{\emptyset, \{0\}, \{1\}, \{0, 1\}\}$
- $\mathcal{P}(B) = ?$

$$\begin{aligned} \mathcal{P}(B) &= \mathcal{P}(A) \cup \left( \bigcup_{s \in \mathcal{P}(A)} \{s \cup \{2\}\} \right) \\ &= \{\emptyset, \{0\}, \{1\}, \{0, 1\}\} \cup \{\{2\}, \{0, 2\}, \{1, 2\}, \{0, 1, 2\}\} \\ &= \{\emptyset, \{0\}, \{1\}, \{0, 1\}, \{2\}, \{0, 2\}, \{1, 2\}, \{0, 1, 2\}\} \end{aligned}$$



# Cardinality of a Power Set: $|\mathcal{P}(S)| = 2^{|S|}$

- $A = \{0, 1\}$  and  $B = \{0, 1, 2\}$
- $\mathcal{P}(A) = \{\emptyset, \{0\}, \{1\}, \{0, 1\}\}$
- $\mathcal{P}(B) = \{\emptyset, \{0\}, \{1\}, \{2\}, \{0, 1\}, \{0, 2\}, \{1, 2\}, \{0, 1, 2\}\}$





# Cardinality of a Power Set: $|\mathcal{P}(S)| = 2^{|S|}$

- $A = \{0, 1\}$  and  $B = \{0, 1, 2\}$
- $\mathcal{P}(A) = \{\emptyset, \{0\}, \{1\}, \{0, 1\}\}$
- $\mathcal{P}(B) = \{\emptyset, \{0\}, \{1\}, \{2\}, \{0, 1\}, \{0, 2\}, \{1, 2\}, \{0, 1, 2\}\}$
- $|\mathcal{P}(B)| = ?$

$$\begin{aligned}
 |\mathcal{P}(B)| &= |\mathcal{P}(A)| + \left| \bigcup_{s \in \mathcal{P}(A)} \{s \cup \{2\}\} \right| \\
 &= |\mathcal{P}(A)| + \sum_{s \in \mathcal{P}(A)} |\{s \cup \{2\}\}| \\
 &= |\mathcal{P}(A)| + |\mathcal{P}(A)| \\
 &= 2 \cdot |\mathcal{P}(A)|
 \end{aligned}$$



# Cardinality of a Power Set: $|\mathcal{P}(S)| = 2^{|S|}$

- $A = \{0, 1\}$  and  $B = \{0, 1, 2\}$
- $\mathcal{P}(A) = \{\emptyset, \{0\}, \{1\}, \{0, 1\}\}$
- $\mathcal{P}(B) = \{\emptyset, \{0\}, \{1\}, \{2\}, \{0, 1\}, \{0, 2\}, \{1, 2\}, \{0, 1, 2\}\}$
- $|\mathcal{P}(B)| = 2 \cdot |\mathcal{P}(A)| = 2 \cdot 2^{|A|} = 2^{|A|+1} = 2^{|B|}$



# Relations

## Definition (Relation)

A *relation* between two sets is a subset of their Cartesian product.



# Relations

## Definition (Relation)

A *relation* between two sets is a subset of their Cartesian product.

Given  $A = \{a, b, c\}$  and  $B = \{0, 1, 2\}$ :

$$A \times B = \{(a, 0), (a, 1), (a, 2), (b, 0), (b, 1), (b, 2), (c, 0), (c, 1), (c, 2)\}$$



# Relations

## Definition (Relation)

A *relation* between two sets is a subset of their Cartesian product.

Given  $A = \{a, b, c\}$  and  $B = \{0, 1, 2\}$ :

$$A \times B = \{(a, 0), (a, 1), (a, 2), (b, 0), (b, 1), (b, 2), (c, 0), (c, 1), (c, 2)\}$$

A sample relation might be:

$$\mathcal{R} = \{(a, 0), (a, 1), (a, 2), (b, 1), (b, 2), (c, 2)\}$$



# Relations

## Definition (Relation)

A *relation* between two sets is a subset of their Cartesian product.

Given:

$$A \times A = \{(a, a), (a, b), (a, c), (b, a), (b, b), (b, c), (c, a), (c, b), (c, c)\}$$



# Relations

## Definition (Relation)

A *relation* between two sets is a subset of their Cartesian product.

Given:

$$A \times A = \{(a, a), (a, b), (a, c), (b, a), (b, b), (b, c), (c, a), (c, b), (c, c)\}$$

A sample relation might be:

$$\mathcal{O} = \{(a, b), (a, c), (b, c)\}$$



# Equivalence Relation

## Definition (Equivalence Relation)

A relation between a set and its self is an *equivalence relation* if and only if it is *reflexive*, *symmetric*, and *transitive*.





# Equivalence Relation

## Definition (Equivalence Relation)

A relation between a set and its self is an *equivalence relation* if and only if it is *reflexive*, *symmetric*, and *transitive*.

Given the relation on  $A$ :

$$\mathcal{O} = \{(a, b), (a, c), (b, c)\}$$



# Equivalence Relation

## Definition (Equivalence Relation)

A relation between a set and its self is an *equivalence relation* if and only if it is *reflexive*, *symmetric*, and *transitive*.

Given the relation on  $A$ :

$$\mathcal{O} = \{(a, b), (a, c), (b, c)\}$$

Since  $a$  does not relate to its self ( $a \not\sim a$ ) this is not *reflexive*.



# Equivalence Relation

## Definition (Equivalence Relation)

A relation between a set and its self is an *equivalence relation* if and only if it is *reflexive*, *symmetric*, and *transitive*.

Given the relation on  $A$ :

$$\mathcal{O} = \{(a, b), (a, c), (b, c)\}$$

Since  $a$  relates to  $b$  ( $a \sim b$ ) but  $b$  does not relate to  $a$  ( $b \not\sim a$ ) this is not *symmetric*.



# Equivalence Relation

## Definition (Equivalence Relation)

A relation between a set and its self is an *equivalence relation* if and only if it is *reflexive*, *symmetric*, and *transitive*.

Given the relation on  $A$ :

$$\mathcal{O} = \{(a, b), (a, c), (b, c)\}$$

Since  $a \sim b$  and  $b \sim c$  and  $a \sim c$  this is *transitive*.



# Equivalence Relation

## Definition (Equivalence Relation)

A relation between a set and its self is an *equivalence relation* if and only if it is *reflexive*, *symmetric*, and *transitive*.

Given the relation on  $A$ :

$$\mathcal{C} = \{(a, a), (a, b), (b, a), (b, b), (c, c)\}$$



# Equivalence Relation

## Definition (Equivalence Relation)

A relation between a set and its self is an *equivalence relation* if and only if it is *reflexive*, *symmetric*, and *transitive*.

Given the relation on  $A$ :

$$\mathcal{C} = \{(a, a), (a, b), (b, a), (b, b), (c, c)\}$$

Since  $a \sim a$ ,  $b \sim b$ , and  $c \sim c$  this is *reflexive*.



# Equivalence Relation

## Definition (Equivalence Relation)

A relation between a set and its self is an *equivalence relation* if and only if it is *reflexive*, *symmetric*, and *transitive*.

Given the relation on  $A$ :

$$\mathcal{C} = \{(a, a), (a, b), (b, a), (b, b), (c, c)\}$$

Since  $a \sim b$  and  $b \sim a$  this is *symmetric*.



# Equivalence Relation

## Definition (Equivalence Relation)

A relation between a set and its self is an *equivalence relation* if and only if it is *reflexive*, *symmetric*, and *transitive*.

Given the relation on  $A$ :

$$\mathcal{C} = \{(a, a), (a, b), (b, a), (b, b), (c, c)\}$$

Since  $a \sim b$ ,  $b \sim a$  and  $a \sim a$  (also,  $b \sim a$ ,  $a \sim b$ , and  $b \sim b$ ) this is *transitive*.





# Equivalence Relation

## Definition (Equivalence Relation)

A relation between a set and its self is an *equivalence relation* if and only if it is *reflexive*, *symmetric*, and *transitive*.

Given the relation on  $A$ :

$$\mathcal{C} = \{(a, a), (a, b), (b, a), (b, b), (c, c)\}$$

This relation is an *equivalence relation*.




# Equivalence Relation

## Definition (Equivalence Relation)

A relation between a set and its self is an *equivalence relation* if and only if it is *reflexive*, *symmetric*, and *transitive*.

Given the relation on  $A$ :

$$\mathcal{C} = \{(a, a), (a, b), (b, a), (b, b), (c, c)\}$$



# Function

## Definition (Function)

A *function* is a relation between two sets, the first called the *domain* and the second the *co-domain*, such that for all  $x$  in the domain there exists a unique  $y$  in the co-domain such that  $(x, y)$  is in the relation.



# Function

## Definition (Function)

A *function* is a relation between two sets, the first called the *domain* and the second the *co-domain*, such that for all  $x$  in the domain there exists a unique  $y$  in the co-domain such that  $(x, y)$  is in the relation.

Given:

$$A \times B = \{(a, 0), (a, 1), (a, 2), (b, 0), (b, 1), (b, 2), (c, 0), (c, 1), (c, 2)\}$$

The relation:

$$\mathcal{R} = \{(a, 0), (a, 1), (a, 2), (b, 1), (b, 2), (c, 2)\}$$

is not a function



# Function

## Definition (Function)

A *function* is a relation between two sets, the first called the *domain* and the second the *co-domain*, such that for all  $x$  in the domain there exists a unique  $y$  in the co-domain such that  $(x, y)$  is in the relation.

Given:

$$A \times B = \{(a, 0), (a, 1), (a, 2), (b, 0), (b, 1), (b, 2), (c, 0), (c, 1), (c, 2)\}$$

But, the relation:


$$S = \{(a, 1), (b, 2), (c, 0)\}$$

is a function



# Visualizing Functions

Domain    Co-Domain




1 - 1 & onto




# Visualizing Functions

Domain Co-Domain



1 - 1 & onto

Domain Co-Domain




onto




# Visualizing Functions

Domain Co-Domain




1 - 1 & onto

Domain Co-Domain



1 - 1

Domain Co-Domain




onto






# Visualizing Functions

Domain Co-Domain




1 - 1 & onto

Domain Co-Domain




1 - 1

Domain Co-Domain




onto

Domain Co-Domain




# Visualizing Functions

Domain Co-Domain




1 - 1 & onto

Domain Co-Domain




1 - 1

Domain Co-Domain




Not a function

Domain Co-Domain



onto

Domain Co-Domain




# Table of Contents


- 1 Sets, Relations, and Functions
- 2 Graph Theory**
- 3 Theorems and Proofs
- 4 Next Class



## Graphs



# Graphs




- Vertex Set:

$$V = \{A, B, C, D\}$$



# Graphs



- Vertex Set:


$$V = \{A, B, C, D\}$$

- Edge Set:

$$E = \{e_1, e_2, e_3, e_4, e_5, e_6\}$$



# Graphs



- Vertex Set:

$$V = \{A, B, C, D\}$$

- Edge Set:


$$E = \{e_1, e_2, e_3, e_4, e_5, e_6\}$$

- Edge Set:

$$E = \{(A, A), (A, B), (A, D), (B, B), (B, D), (C, D)\}$$



# Graphs



- Vertex Set:

$$V = \{A, B, C, D\}$$

- Edge Set:

$$E = \{e_1, e_2, e_3, e_4, e_5, e_6\}$$

- Edge Set:

$$E = \{(A, A), (A, B), (A, D), (B, B), (B, D), (C, D)\}$$

- Graph:

$$G = (V, E)$$


$$= (\{A, B, C, D\},$$

$$\{(A, A), (A, B), (A, D), (B, B), (B, D), (C, D)\})$$






# Types of Graphs




# Types of Graphs

## Graph




## Directed Graph




# Types of Graphs


## Graph



## Bipartite Graph




## Directed Graph




# Types of Graphs


## Graph




## Bipartite Graph



## Directed Graph




## Complete Graph




## Types of Graphs


Graph




Bipartite Graph




Tree



Directed Graph




Complete Graph




## Types of Graphs


Graph




Bipartite Graph




Tree




Directed Graph




Complete Graph



Binary Tree




# Relations as Graphs



# Relations as Graphs

Equivalence Relation?







# Relations as Graphs

Equivalence Relation?

- Reflexive ✓



# Relations as Graphs




Equivalence Relation?

- Reflexive ✓
- Symmetric ✓



# Relations as Graphs




Equivalence Relation?

- Reflexive ✓
- Symmetric ✓
- Transitive ✓



# Relations as Graphs




Equivalence Relation?

- Reflexive ✓
- Symmetric ✓
- Transitive ✓



# Relations as Graphs



## Equivalence Relation ✓

- Reflexive ✓
- Symmetric ✓
- Transitive ✓
- Equivalence Classes

$$A = \{a, b, d\} \text{ \& } C = \{c\}$$



# Table of Contents

- 1 Sets, Relations, and Functions
- 2 Graph Theory
- 3 Theorems and Proofs**
- 4 Next Class



# Direct Proof

## Theorem (De Morgan's Law)

*Given two sets  $A$  and  $B$ , the complement of their union is equal to the intersection of their complements:*

$$(A \cup B)^c = A^c \cap B^c.$$



# Direct Proof

## Theorem (De Morgan's Law)

*Given two sets  $A$  and  $B$ , the complement of their union is equal to the intersection of their complements:*

$$(A \cup B)^c = A^c \cap B^c.$$

*Proof:* Let  $A$  and  $B$  be sets and  $x \in (A \cup B)^c$ , thus  $x \notin A \cup B$ .





# Direct Proof

## Theorem (De Morgan's Law)

*Given two sets  $A$  and  $B$ , the complement of their union is equal to the intersection of their complements:*

$$(A \cup B)^c = A^c \cap B^c.$$

*Proof:* Let  $A$  and  $B$  be sets and  $x \in (A \cup B)^c$ , thus  $x \notin A \cup B$ . This means  $x \notin A$  and  $x \notin B$ , so  $x \in A^c$  and  $x \in B^c$ .



# Direct Proof

## Theorem (De Morgan's Law)

*Given two sets  $A$  and  $B$ , the complement of their union is equal to the intersection of their complements:*

$$(A \cup B)^c = A^c \cap B^c.$$

*Proof:* Let  $A$  and  $B$  be sets and  $x \in (A \cup B)^c$ , thus  $x \notin A \cup B$ . This means  $x \notin A$  and  $x \notin B$ , so  $x \in A^c$  and  $x \in B^c$ . By definition then,  $x \in A^c \cap B^c$  and  $(A \cup B)^c \subseteq A^c \cap B^c$ .



# Direct Proof

## Theorem (De Morgan's Law)

*Given two sets  $A$  and  $B$ , the complement of their union is equal to the intersection of their complements:*

$$(A \cup B)^c = A^c \cap B^c.$$

*Proof:* Let  $A$  and  $B$  be sets and  $x \in (A \cup B)^c$ , thus  $x \notin A \cup B$ . This means  $x \notin A$  and  $x \notin B$ , so  $x \in A^c$  and  $x \in B^c$ . By definition then,  $x \in A^c \cap B^c$  and  $(A \cup B)^c \subseteq A^c \cap B^c$ .

Now suppose  $x \in A^c \cap B^c$  or equivalently  $x \in A^c$  and  $x \in B^c$ .



# Direct Proof

## Theorem (De Morgan's Law)

*Given two sets  $A$  and  $B$ , the complement of their union is equal to the intersection of their complements:*

$$(A \cup B)^c = A^c \cap B^c.$$

*Proof:* Let  $A$  and  $B$  be sets and  $x \in (A \cup B)^c$ , thus  $x \notin A \cup B$ . This means  $x \notin A$  and  $x \notin B$ , so  $x \in A^c$  and  $x \in B^c$ . By definition then,  $x \in A^c \cap B^c$  and  $(A \cup B)^c \subseteq A^c \cap B^c$ .

Now suppose  $x \in A^c \cap B^c$  or equivalently  $x \in A^c$  and  $x \in B^c$ . This tells us that  $x \notin A$  and  $x \notin B$  and thus  $x \notin A \cup B$ , i.e.  $x \in (A \cup B)^c$ .



# Direct Proof

## Theorem (De Morgan's Law)

Given two sets  $A$  and  $B$ , the complement of their union is equal to the intersection of their complements:

$$(A \cup B)^c = A^c \cap B^c.$$

*Proof:* Let  $A$  and  $B$  be sets and  $x \in (A \cup B)^c$ , thus  $x \notin A \cup B$ . This means  $x \notin A$  and  $x \notin B$ , so  $x \in A^c$  and  $x \in B^c$ . By definition then,  $x \in A^c \cap B^c$  and  $(A \cup B)^c \subseteq A^c \cap B^c$ .

Now suppose  $x \in A^c \cap B^c$  or equivalently  $x \in A^c$  and  $x \in B^c$ . This tells us that  $x \notin A$  and  $x \notin B$  and thus  $x \notin A \cup B$ , i.e.  $x \in (A \cup B)^c$ . Therefore,  $A^c \cap B^c \subseteq (A \cup B)^c$  and

$$(A \cup B)^c = A^c \cap B^c$$

as desired.



# Direct Proof

## Theorem (De Morgan's Law)

Given two sets  $A$  and  $B$ , the complement of their union is equal to the intersection of their complements:

$$(A \cup B)^c = A^c \cap B^c.$$

*Proof:* Let  $A$  and  $B$  be sets and  $x \in (A \cup B)^c$ , thus  $x \notin A \cup B$ . This means  $x \notin A$  and  $x \notin B$ , so  $x \in A^c$  and  $x \in B^c$ . By definition then,  $x \in A^c \cap B^c$  and  $(A \cup B)^c \subseteq A^c \cap B^c$ .

Now suppose  $x \in A^c \cap B^c$  or equivalently  $x \in A^c$  and  $x \in B^c$ . This tells us that  $x \notin A$  and  $x \notin B$  and thus  $x \notin A \cup B$ , i.e.  $x \in (A \cup B)^c$ . Therefore,  $A^c \cap B^c \subseteq (A \cup B)^c$  and

$$(A \cup B)^c = A^c \cap B^c$$

as desired. Why did this proof require two parts?



# By Cases

## Theorem

*Given any integer  $n$ , either  $n^2$  or  $n^2 - 1$  is divisible by four.*



# By Cases

## Theorem

*Given any integer  $n$ , either  $n^2$  or  $n^2 - 1$  is divisible by four.*

*Proof:* (Case 1) Let  $n$  be an even integer so that we may write  $n = 2k$  for some unique  $k$ . Then

$$n^2 = 4k^2$$

and  $n^2$  is divisible by four.





# By Cases

## Theorem

*Given any integer  $n$ , either  $n^2$  or  $n^2 - 1$  is divisible by four.*

*Proof:* (Case 1) Let  $n$  be an even integer so that we may write  $n = 2k$  for some unique  $k$ . Then

$$n^2 = 4k^2$$

and  $n^2$  is divisible by four.

(Case2) Now, if  $n$  is an odd integer then we write  $n = 2k + 1$  for some unique  $k$ . Thus,

$$n^2 - 1 = 4k^2 + 4k + 1 - 1 = 4(k^2 + k)$$

and  $n^2 - 1$  is divisible by four.



# By Cases

## Theorem

*Given any integer  $n$ , either  $n^2$  or  $n^2 - 1$  is divisible by four.*

*Proof:* (Case 1) Let  $n$  be an even integer so that we may write  $n = 2k$  for some unique  $k$ . Then

$$n^2 = 4k^2$$

and  $n^2$  is divisible by four.

(Case2) Now, if  $n$  is an odd integer then we write  $n = 2k + 1$  for some unique  $k$ . Thus,

$$n^2 - 1 = 4k^2 + 4k + 1 - 1 = 4(k^2 + k)$$

and  $n^2 - 1$  is divisible by four.

Therefore, for any integer  $n$  we have shown that either  $n^2$  or  $n^2 - 1$  is divisible by four.



# Contrapositive

## Theorem

*If  $n^2$  is even, then  $n$  is even.*



# Contrapositive

## Theorem

*If  $n^2$  is even, then  $n$  is even.*

*Proof:* Suppose that  $n$  is odd and is written  $n = 2k + 1$  for some unique  $k$ .



# Contrapositive

## Theorem

*If  $n^2$  is even, then  $n$  is even.*

*Proof:* Suppose that  $n$  is odd and is written  $n = 2k + 1$  for some unique  $k$ . Then we can write

$$n^2 = (2k + 1)^2 = 4k^2 + 4k + 1 = 2(2k^2 + 2k) + 1$$

which is odd.



# Contrapositive

## Theorem

*If  $n^2$  is even, then  $n$  is even.*

*Proof:* Suppose that  $n$  is odd and is written  $n = 2k + 1$  for some unique  $k$ . Then we can write

$$n^2 = (2k + 1)^2 = 4k^2 + 4k + 1 = 2(2k^2 + 2k) + 1$$

which is odd. Therefore, if  $n$  is odd, then  $n^2$  is odd and so if  $n^2$  is even, then  $n$  is even.



# Contradiction

## Theorem

*No integer is both even and odd.*



# Contradiction

## Theorem

*No integer is both even and odd.*

*Proof:* Suppose that  $n$  is both even and odd so that  $n = 2k$  and  $n = 2l + 1$  for some unique  $k$  and  $l$ .





# Contradiction

## Theorem

*No integer is both even and odd.*

*Proof:* Suppose that  $n$  is both even and odd so that  $n = 2k$  and  $n = 2l + 1$  for some unique  $k$  and  $l$ . Then we can write  $2k = 2l + 1$  and  $1 = 2(k - l)$ .



# Contradiction

## Theorem

*No integer is both even and odd.*

*Proof:* Suppose that  $n$  is both even and odd so that  $n = 2k$  and  $n = 2l + 1$  for some unique  $k$  and  $l$ . Then we can write  $2k = 2l + 1$  and  $1 = 2(k - l)$ . If  $k - l = 0$ , then  $1 = 0$  and if  $k - l \neq 0$ , then 2 divides 1.



# Contradiction

## Theorem

*No integer is both even and odd.*

*Proof:* Suppose that  $n$  is both even and odd so that  $n = 2k$  and  $n = 2l + 1$  for some unique  $k$  and  $l$ . Then we can write  $2k = 2l + 1$  and  $1 = 2(k - l)$ . If  $k - l = 0$ , then  $1 = 0$  and if  $k - l \neq 0$ , then 2 divides 1. In either case we derive a contradiction and therefore no integer is both even and odd.



# Induction

## Theorem

*Any tree with  $n$  vertices has  $n - 1$  edges.*



# Induction

## Theorem

*Any tree with  $n$  vertices has  $n - 1$  edges.*



(Base Case) When there is only one vertex there are no edges since trees do not contain loops and there is not a second vertex to connect to.




# Induction

## Theorem

*Any tree with  $n$  vertices has  $n - 1$  edges.*


(Induction Step) Assume that the theorem is true for some  $k \geq 2$  and consider a tree with  $k + 1 \geq 3$  vertices.



# Induction

## Theorem

*Any tree with  $n$  vertices has  $n - 1$  edges.*




(Induction Step) Assume that the theorem is true for some  $k \geq 2$  and consider a tree with  $k + 1 \geq 3$  vertices. Since there are at least two vertices the tree must contain at least one leaf.



# Induction

## Theorem

*Any tree with  $n$  vertices has  $n - 1$  edges.*



(Induction Step) Assume that the theorem is true for some  $k \geq 2$  and consider a tree with  $k + 1 \geq 3$  vertices. Since there are at least two vertices the tree must contain at least one leaf. Removing the leaf removes one vertex and one edge; we now have a subtree with  $k$  vertices and, by induction,  $k - 1$  edges.






# Induction

## Theorem

*Any tree with  $n$  vertices has  $n - 1$  edges.*



(Induction Step) Assume that the theorem is true for some  $k \geq 2$  and consider a tree with  $k + 1 \geq 3$  vertices. Since there are at least two vertices the tree must contain at least one leaf. Removing the leaf removes one vertex and one edge; we now have a subtree with  $k$  vertices and, by induction,  $k - 1$  edges. Thus the original tree has  $k + 1$  vertices and  $k$  edges.



# Table of Contents

- 1 Sets, Relations, and Functions
- 2 Graph Theory
- 3 Theorems and Proofs
- 4 Next Class



# Next Class

- Deterministic and Non-Deterministic Finite State Automata



# Discrete Math Review

Dr. Chuck Rocca  
roccac@wcsu.edu

<http://sites.wcsu.edu/roccac>

